

POSTER PRESENTATION

Open Access

Muscle wasting in icu trauma patients: quantitative and qualitative changes

MG Annetta^{1*}, D Silvestri¹, DL Grieco¹, MF La Torre², N Magarelli², C Policola³, A Caricato¹, S Della Casa³, M Antonelli¹

From ESICM LIVES 2015

Berlin, Germany. 3-7 October 2015

Introduction

Loss of skeletal muscle is a typical and early finding in critically ill patients, mostly related to forced immobility, systemic inflammatory response, and protein depletion; it is clinically associated with a functional impairment that may cause difficult weaning from mechanical ventilation, increased length of ICU and hospital stay and long term functional disability.

Objectives

We carried out a prospective observational study to evaluate the morphological changes of the muscle mass in a group of enterally fed trauma patients, comparing them with a control group of malnourished patients (anorexia nervosa).

Methods

Fifteen ICU patients were studied on day 1 and on day 20 after trauma and compared with 15 anorexic patients with severe malnutrition (BMI < 17) volunteering to participate to the study. Muscle mass was evaluated by ultrasound scan, a recently validated method with several advantages: it is accurate, non-invasive, inexpensive, easily repeatable, easy to perform at bedside and apt to detect both qualitative and quantitative changes of the muscle. In all patients, we studied the rectus femoris (RF) and anterior tibialis (AT) muscles, considering

- the anterior-posterior diameter,
- the lateral diameter,
- the sectional area and
- the muscle echogenicity according to Heckmatt's score.

Results

Our trauma patients were 12 males and 3 females; age was 53 (range 43-61), length of ICU stay 35 days (23-50), days on mechanical ventilation 22 (12-36), injury severity score 34 (27-50), BMI 27.5 (23.1-29.4). The anorexic patients were all female: age 21.5 (17-30), BMI 14.9 (13.2-16.65). From day 1 to day 20 after trauma, the muscle mass decreased significantly, reaching the values observed in the anorexic patients: the anterior-posterior diameter of both RF and AT decreased significantly ($p < 0.008$ and $p < 0.001$, respectively); the area of RF decreased by 26.1% (16.1% - 36.3%, $p < 0.001$); the area of AT decreased by 37.3% (24.5% - 50.2%, $p < 0.001$), reaching a value not dissimilar from anorexic patients ($p = 0.18$). On day 1 after trauma, muscle echogenicity of both RF and TA was higher than in anorexic patients ($p < 0.001$), suggesting very early inflammatory changes. The muscle echogenicity was higher on day 20 if compared to day 1, both in RF ($p < 0.005$) and in AT ($p < 0.004$), suggesting further abnormalities of muscle structure which cannot be explained by malnutrition alone.

Conclusions

Evaluation of skeletal muscle changes in the critically ill patients can be successfully performed by ultrasound, which yields both quantitative and qualitative data. Our results confirm the progressive depletion of muscle mass in the ICU trauma patient; also, our data suggest that the inflammatory process - which is a major contributor of muscle wasting - begins very early in the critical illness, in the first 24 hrs after trauma, and progress further during the ICU stay.

¹Catholic University of Rome, Anesthesiology and Intensive Care Medicine, Rome, Italy

Full list of author information is available at the end of the article

Authors' details

¹Catholic University of Rome, Anesthesiology and Intensive Care Medicine, Rome, Italy. ²Catholic University of Rome, Diagnostic Radiology, Rome, Italy. ³Catholic University of Rome, Endocrinology, Rome, Italy.

Published: 1 October 2015

doi:10.1186/2197-425X-3-S1-A181

Cite this article as: Annetta et al.: Muscle wasting in icu trauma patients: quantitative and qualitative changes. *Intensive Care Medicine Experimental* 2015 **3**(Suppl 1):A181.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
